

New theorems in approximation theory

Nabaa Mery Kasim*

Date of acceptance 12/5 / 2009

Abstract:

The aim of this paper is to prove some results for equivalence of moduli of smoothness in approximation theory , we used a"non uniform" modulus of smoothness $\varpi_{\phi}^k(f,t) = \sup_{0 < h \leq t} \left\| \Delta_{\phi}^k(f,x) \right\|_p^{-k}$ and the weighted Ditzian –Totik moduli of smoothness $\varpi_{k,v}^{\psi}(f,t) = \sup_{0 < h \leq t} \left\| \psi(\cdot)^v \Delta_{h\psi(\cdot)}^k(f,x) \right\|_p^{-k}$ in $L_p[-1,1], 0 < p \leq \infty$ by spline functions ,several results are obtained .For example , it shown that ,for any $1 \leq \phi \leq r+1, 1 \leq p \leq \infty$ the inequality $|J|^v \varpi_{\phi-v}^k(S^{(v)}, J)_p \approx \varpi_{\phi}^k(S, J)_p$, $1 \leq v \leq \min[\phi, m+1]$ is satisfied ,finally, similar result for chebyshev partition and weighted Ditzian –Totik moduli of smoothness are also obtained.

Key words: Univariate splines , moduli of smoothness , "non uniform" modulus of smoothness , weighted Ditzian –Totik moduli of smoothness.

Introduction:

Equivalence moduli of smoothness was proved by Hu and Yu [1] and by Hu [2] for uniform and quasi-uniform partition , respectively. In fact , if we set $m = k - 1$, then corollary (1.5) (with an additional restriction $k \leq r$) becomes theorem 1 in [2] ,also in the case $k = r + 1$ and $m = r - 1$, corollary (1.5) follow from theorem 2 of [2] .

A function S is called a spline if : [3]

- 1- the domain is an interval $[a,b]$.
- 2- $S, S', S'', \dots, S^{(r-1)}$ are all continuous function on $[a,b]$
- 3- there are point (the knot of S)such that $a < Z_0 < Z_1 < \dots < Z_{n-1} < Z_n = b$ and such that S is a polynomial of

degree $\leq r$ on each subinterval $[Z_i, Z_{i+1}]$

Usually piecewise polynomials from $S_r(Z_n)$ are called "splines" if they possess continuous $(r-1)st$ derivatives .We emphasize that if $S \in S_r(Z_n)$ and $m \in N$ then $S^{(m)} \in \delta_{\max\{r-m,0\}}(Z_n)$ and $S^{(m)} = 0$ a.e if $m \geq r+1$.

Now a moduli of smoothness are intended for mathematicians working in approximation theory ,numerical analysis and real analysis .Measuring the smoothness of function by differentiability is too crude for many purposes in approximation theory . more suitable measurement are provided by moduli of smoothness.

*Math .Department/College of Education /Abn-Al Haitham/ University of Baghdad
[4]

$L_p(J), 0 < p \leq \infty$, denotes the space of all measurable function f on J . Such that $\|f\|_{L_p(J)} < \infty$.

The k th symmetric of f is given by

$$\Delta_\beta^k(f, x, [a, b]) = \begin{cases} \sum_{i=0}^k (-1)^{k-i} \binom{k}{i} f(x + i\beta) & \text{if } [x, x+k\beta] \subset [a, b] \\ 0 & \text{otherwise} \end{cases}$$

then the k th "non uniform" modulus of smoothness of $f \in L_p[-1, 1], 0 < p \leq \infty$ is define by

$$\varpi_\phi^k(f, t) = \sup_{0 < h \leq t} \left\| \Delta_\beta^k(f, x) \right\|_p$$

Where

$$\beta = \beta(x, h) = \sqrt{1 - x^2} h + h^2, \phi(x) = \sqrt{1 - x^2}$$

and

The weighted Ditzian –Totik moduli of smoothness of a function $f \in L_p[-1, 1], 0 < p \leq \infty$ is define by

$$\varpi_{k,v}^\psi(f, t) = \sup_{0 < h \leq t} \left\| \psi(\cdot)^v \Delta_{h\psi(\cdot)}^k(f, x) \right\|_p, [5]$$

$$\text{Where } \psi(x) = \sqrt{1 - x^2}$$

A partition $Z_n = \{Z_0, \dots, Z_n\} = -1 < Z_0 < Z_1 < \dots < Z_{n-1} < Z_n = 1$ of the interval $[-1, 1]$ denote the scale of parition Z_n by $\theta = \theta(Z_n) = \max \frac{|J_i|}{|J_j|}$ where $J_j = [Z_j, Z_{j+1}]$ with $Z_j = -1, j < 0$ and $Z_j = 1, j > n$.

Auxiliary results

The following properties of the moduli of smoothness are well known[6],[7]

(i) for

$f \in L_p(J), 0 < p \leq \infty$, we have

$$\omega_{k+1}(f, t, J)_p \leq 2^{\max\{1, \frac{1}{p}\}} \omega_k(f, t, J)_p, k \in N \quad (2.1)$$

(ii)

$$\omega_k(f, \lambda t, J)_p \leq C(k, p)(\lambda + 1)^{k-1+\max\{1, \frac{1}{p}\}} \omega_k(f, t, J)_p, \lambda > 0 \quad (2.2)$$

(iii) suppose that $f \in L_p[-1, 1], 0 < p < \infty$ and $k, \mu \in N$ then

$$\sum_{j=0}^{n-\mu-1} \omega_k(f, \bigcup_{i=j}^{j+\mu} J_i)_p^p \leq \begin{cases} c(k, \Delta, \mu, p) \omega_k(f, n^{-1})_p^p & \text{if } z_n = u_n^\Delta \\ c(k, \mu, p) \omega_k^\phi(f, n^{-1})_p^p & \text{if } z_n = t_n \end{cases} \quad (2.3)$$

Lemma .1 (Whitney's inequality)

). [8]

For any $f \in L_p[a, b], 0 < p \leq \infty$ such that $q_{k-1} \in \Pi_{k-1}$ such that

$$\|f - q_{k-1}\|_{L_p[a, b]} \leq C \omega_k(f, [a, b]).$$

(2.4)

Lemma .2

For any polynomial $q_k \in \Pi_k, 0 < p \leq \infty$ and intervals I and J such that $I \subseteq J$ we have

$$\begin{aligned} |J|^{\frac{1}{p}} \|q_r^{(v)}\|_{L_\infty(J)} &\approx \|q_r^{(v)}\|_{L_p(J)} \leq \\ C(r, \frac{|J|}{|I|}, p) |J|^{-v} \|q_r\|_{L_p(I)}, 0 \leq v \leq r & \end{aligned} \quad (2.5)$$

The main result

The following theorem is not satisfied for all $f \in L_p, 0 < p < \infty$ but satisfied for spline functions

Theorem .1 (local estimates)

Let $S \in S_r(Z_n), r \in N$ and

$J = [Z_{M_1}, Z_{M_2}]$ with $M_2 - M_1 \leq c_0$ for some constant c_0

Then for $1 \leq \phi \leq r+1, 0 \leq p \leq \infty$ we have

$$\varpi_{\phi^{-v}}^k(S^{(v)}, J)_p \leq c(r, \theta, c_0, p) |J|^{-v} \varpi_\phi^k(S, J)_p$$

for all $v = 1, 2, \dots, \phi$.

proof:

Every where in this proof q_{k-1} denote a polynomial of degree $\leq k-1$ $\exists \|s - q_{k-1}\|_{L_p(J)} \leq C \varpi_\phi^k(s, J)_p$ (2.4)

for any $1 \leq v \leq \phi$. denoting $s_j = \frac{s}{J_j}$ and using [(2.5),[7]] we have

$$|\Delta_h^{k-v}(s^{(v)}, x, J)| \leq |\tilde{\Delta}_p^{k-v}(s^{(v)}, x, J)| \text{ since}$$

$$\varpi_\phi^k(f, t) \approx \omega_\phi^k(f, t)_\infty$$

$$= |\tilde{\Delta}_p^{k-v}(s^{(v)} - q_{k-1}^{(v)}, x, J)|$$

$$\leq 2^{k-v} \|s^{(v)} - q_{k-1}^{(v)}\|_{L_\infty(J)}$$

$$\leq C(k) \max_{M_1 \leq j \leq M_2-1} \|s^{(v)} - q_{k-1}^{(v)}\|_{L_\infty(J_j)}$$

$$\leq C(r, p) \max |J_j|^{-v-\frac{1}{p}} \|s_j - q_{k-1}\|_{L_p(J_j)}$$

$$\leq C(r, \theta, c_0, p) |J_j|^{-v-\frac{1}{p}} \|s - q_{k-1}\|_{L_p(J_j)}$$

$$\leq C(r, \theta, c_0, p) |J_j|^{-v-\frac{1}{p}} \varpi_\phi^k(s, J)_p.$$

Corollary .2 (equivalaence of moduli of smoothness)

Let $S \in S_r(Z_n) \cap C^m[-1,1]$, $r \in N$ and $J = [Z_{M_1}, Z_{M_2}]$ with $M_2 - M_1 \leq c_0$ for some constant c_0

Then for $1 \leq \phi \leq r+1$, $1 \leq p \leq \infty$ we have $|J|^v \varpi_{\phi^{-v}}^k(S^{(v)}, J)_p \approx \varpi_\phi^k(S, J)_p$, $1 \leq v \leq \min[\phi, m+1]$

Equivalaence constants above depend only on r, θ, c_0 .

Suppose

that $\delta_{\max} = \delta_{\max}(Z_n) = \max_{0 \leq j \leq n-1} |J_j|$ and $\delta_{\min}(Z_n) = \min_{0 \leq j \leq n-1} |J_j|$ we say that Z_n is Δ -quasi uniform if $\Delta = \delta_{\max}/\delta_{\min}$ is bounded by a constant independent of n , and denote such partition by u_n^Δ , if $Z_n = u_n^\Delta$ then $\frac{2}{n\Delta} \leq \delta_{\min} \leq \frac{2}{n} < \delta_{\max} < \frac{2\Delta}{n}$ and $\theta(Z_n) \leq \Delta$

Therefore $\delta_{\min} \approx \delta_{\max} \approx n^{-1}$ with Equivalaence constants depending only on Δ .

Theorem.3 (quasi uniform partition)

Let u_n^Δ , $n \in N$ be a quasi uniform partition of $[-1,1]$ and let $S \in S_r(u_n^\Delta)$, $r \in N$ then for any $1 \leq \phi \leq r+1$, $0 < p \leq \infty$ we have $\varpi_{\phi^{-v}}^k(S^{(v)}, n^{-1})_p \leq c(r, \Delta, p) n^v \varpi_\phi^k(S, n^{-1})_p$ for all $v = 1, 2, \dots, \phi$.

proof

Since u_n^Δ is a Δ - quasi uniform $\delta_{\max} \approx \delta_{\min} \approx |J_i| \approx n^{-1}$ for all $0 \leq j \leq n-1$ with equivalence constants depending only Δ . Using (2.2) we have

$$\varpi_{\phi^{-v}}^k(S^{(v)}, n^{-1})_p = \varpi_{\phi^{-v}}^k(s^{(v)}, \frac{\phi}{n\delta_{\min}}, \frac{\delta_{\min}}{\phi})$$

$$\leq C(\phi, \Delta, p) \varpi_{\phi^{-v}}^k(s^{(v)}, \frac{\delta_{\min}}{\phi})_p$$

Now, for any $0 < \beta < \frac{\delta_{\min}}{\phi}$ and $x \in J_j$, $0 \leq j \leq n-1$ such that $\Delta_\beta^{\phi-v}(s^{(v)}, x) \neq 0$

All point
 $x - (\phi - v)\beta / \phi^2 + i\beta$, $0 \leq i \leq \phi - v$ are either in $J_{j-1} \bigcup J_j$ or $J_j \bigcup J_{j+1}$

$$\begin{aligned} \varpi_{\phi-v}^k(s^{(v)}, \delta_{\min} / \phi)_p^p &= \sup_{0 \leq \beta \leq \delta_{\min} / \phi} \left\| \Delta_\beta^k(\phi-v)(s^{(v)}, \cdot) \right\|_p^p \\ &= \sup_{0 \leq \beta \leq \delta_{\min} / \phi} \left\| \sum_{j=0}^{n-1} \left\| \Delta_\beta^k(\phi-v)(s^{(v)}, \cdot) \right\|_p^p \right\|_p^p \\ &\leq 2 \\ &\sup_{0 \leq \beta \leq \delta_{\min} / \phi} \sum_{j=0}^{n-1} \left\| \Delta_\beta^k(\phi-v)(s^{(v)}, \cdot, J_{j-1} \bigcup J_j) \right\|_{L_p(J_{j-1} \bigcup J_j)}^p \\ &\leq 2 \sum_{j=1}^{n-1} \varpi_{\phi-v}^k(s^{(v)}, J_{j-1} \bigcup J_j)_p^p \\ &\leq C(r, \Delta, p) \sum_{j=1}^{n-1} |J_{j-1} \bigcup J_j|^{-vp} \varpi_{\phi}^k(s, J_{j-1} \bigcup J_j)_p^p \\ &\leq C(r, \Delta, p) n^{pv} \sum_{j=1}^{n-1} \varpi_{\phi}^k(s, J_{j-1} \bigcup J_j)_p^p \\ &\leq C(r, \Delta, p) n^{pv} \varpi_{\phi}^k(s, n^{-1})_p^p. \end{aligned}$$

Where the last inequality follow from (2.3)

Corollary .4

Let u_n^Δ , $n \in N$ be a quasi uniform partition of $[-1,1]$ and let $S \in S_r(u_n^\Delta) \cap C^m[-1,1]$, $r \in N$, $0 \leq m \leq r-1$ then for any $1 \leq \phi \leq r+1$, $1 \leq p \leq \infty$ we have

$$n^{-v} \varpi_{\phi-v}^k(S^{(v)}, n^{-1})_p \approx \varpi_{\phi}^k(S, n^{-1})_p, \quad 1 \leq v \leq \min[\phi, m+1]$$

Equivalaence constants above depend only on r, Δ .

Let Z_n is a chebyshev partition , if $Z_n = t_n = (t_i)_{i=0}^n$ where $t_i = \cos\left(\frac{(n-i)\Pi}{n}\right)$, $0 \leq i \leq n$.

Theorem.5 (chebyshev knot)

Let t_n be a chebyshev partition of $[-1,1]$ if $s \in S_r(t_n)$, $r \in N$ then , for any $1 \leq \psi \leq r+1$, $0 < p < 1$ we have $\varpi_{\psi-v}^k(S^{(v)}, n^{-1})_p \leq c(r, p) n^v \varpi_{\psi}^k(S, n^{-1})_p$ for all $1 \leq v \leq \psi$.

Proof:

Chebyshev knot: let $J_{j,\phi-v} = [t_{j-3(\psi-v)}, t_{j+4+3(\psi-v)}]$, denote $J_{j,\psi-v}$, and , for each $0 \leq j \leq n-1$ Let $q_j \in \Pi_{\phi-v-1}$ be such that $\|s^{(v)} - q_j\|_{L_p(J_j)} \leq C \varpi_{\phi-v}(s^{(v)}, J_j)_p$ [6], and we assume that $q_j = 0$ if $v = \psi$. Then using theorem 3.1 and the inequality $\psi(x) \leq n|J_j|$, $x \in J_j$, we have

$$\begin{aligned} \omega_{k-v,v}^{\psi}(s^{(v)}, n^{-1})_p &= \sup_{0 < h < n^{-1}} \left\| \psi^v \Delta_h^k(s^{(v)}, \cdot) \right\|_p \\ &= \sup_{0 < h < n^{-1}} \sum_{j=0}^{n-1} \left\| \psi^v \Delta_h^k(s^{(v)}, \cdot) \right\|_{L_p(J_j)} \\ &\leq \sum_{j=0}^{n-1} n^{vp} |J_j|^{-vp} \\ &\sup_{0 < h < n^{-1}} \left\| \psi^v \Delta_h^k(s^{(v)}, \cdot) - q_j \right\|_{L_p(J_j)} \\ &\leq C(k, p) n^{vp} \sum_{j=0}^{n-1} |J_j|^{-vp} \|s^{(v)} - q_j\|_{L_p(J_j)} \\ &\leq C(k, p) n^{vp} \sum_{j=0}^{n-1} |J_j|^{-vp} \omega_{k-v,v}(s^{(v)}, J_j)_p \\ &\leq C(r, p) n^{vp} \sum_{j=0}^{n-1} |J_j|^{-vp} \varpi_{\phi-v}(s^{(v)}, J_j)_p \\ &\leq C(r, p) n^{vp} \varpi_{\psi}^k(s, n^{-1})_p \end{aligned}$$

where the last inequality follows from (2.3)

Corollary.6

Let $s \in S_r(t_n) \cap C^m[-1,1]$
 $r \in N, 0 \leq m \leq r-1$ then , for any
 $1 \leq \psi \leq r+1, 0 < p < 1$ we have
 $n^{-v} \varpi_{\psi-v}^k(S^{(v)}, n^{-1})_p \approx \varpi_{\psi}^k(S, n^{-1})_p$ for
all $1 \leq v \leq \min\{\psi, m+1\}$.

Equivalence constants above depend only on r .

References:

- [1] Hu.Y and Yu.X.M.1995.Discrete modulus of smoothness of splines with equally spaced knots .SIAMJ.Numer .Anal.(32):1428-1435.
- [2] Hu.Y.1999.Some theorems of the moduli of smoothness in approximation theory.J.Aprox.282-293.
- [3] Beatsone. R.K. 1981.Convex approximation by spline . SIAMJ. Math.(12):549-559.
- [4] DeVore.R.A. and Totik.V.1987. Moduli of smoothness.New york .Sprinnger-Verlag.(3):201-215.
- [5] Kopotun.K,Leviatan .D and Shevchuk .2006.Coconvex approximation in the uniform norm,Acta.mat.Hungar.110(1-2):117-151.
- [6] DeVore.R.A and Lorentz.G.G.2003. Constructive approximation . Berlin .springer-verlag,vol.(4): 303-307.
- [7] Petrushev.P.P and Popov.V.A. 1987.Rational approximation of real functions. Cambridge University press.vol.28(5):1231-1236.
- [8] Kopotun.K.A. 2001. Whitney theorem of interpolatory type for K-monotone functions .constructive approximation .17:307-317.

ميرهنا جديدة في نظرية التقرير

*نبأ ميري قاسم

*قسم الرياضيات / كلية التربية / ابن الهيثم / جامعة بغداد

الخلاصة

الهدف من هذا البحث هو برهان نتائج للتفاوتات بين مقاييس النعومة في نظرية التقرير ، استخدمنا مقاييس

$$\text{النعومة غير الطبيعي } \varpi_{\phi}^k(f, t) = \sup_{0 < h \leq t} \left\| \Delta_{\beta}^{-k}(f, x) \right\|_p \text{ ومقاييس نعومة ديتزين الثقيل}$$

$$\text{لدوال السيلاني ، نتائج } L_p[-1,1], 0 < p \leq \infty \text{ في الفضاءات } \varpi_{k,v}^{\psi}(f, t) = \sup_{0 < h \leq t} \left\| \psi(\cdot)^v \Delta_{h\psi(\cdot)}^{-k}(f, x) \right\|_p$$

عديدة حصلنا عليها مثلا، بينما المتراجحة $|J|^v \varpi_{\phi-v}^k(S^{(v)}, J)_p \approx \varpi_{\phi}^k(S, J)_p$ عندما $1 \leq v \leq \min[\phi, m+1]$ ، $1 \leq \phi \leq r+1$, $1 \leq p \leq \infty$ متحققة واخيرا بينما تجزئة شيبشيف متحققة ايضا باستخدام مقاييس نعومة ديتزين الثقيل.