

2025

k –Mixed Modulus of Smoothness

Rehab Amer Kamel

Department of Mathematics, College of Education for Pure Sciences, University of Babylon, Hillah, Iraq,
pure.rehab.amer@uobabylon.edu.iq

Eman Samir Bhaya

Department of Mathematics, College of Education, Al-Zahraa University for Women, Karbala, Iraq AND
Department of Mathematics, College of Education for Pure Sciences, University of Babylon, Hillah, Iraq,
emanbhaya@alzahraa.edu.iq

Follow this and additional works at: <https://bsj.researchcommons.org/home>

How to Cite this Article

Kamel, Rehab Amer and Bhaya, Eman Samir (2025) "k –Mixed Modulus of Smoothness," *Baghdad Science Journal*: Vol. 22: Iss. 1, Article 21.

DOI: 10.21123/bsj.2024.9726

Available at: <https://bsj.researchcommons.org/home/vol22/iss1/21>

This Article is brought to you for free and open access by Baghdad Science Journal. It has been accepted for inclusion in Baghdad Science Journal by an authorized editor of Baghdad Science Journal.



RESEARCH ARTICLE

$\mathbb{k}$ -Mixed Modulus of Smoothness

Rehab Amer Kamel^{1,*}, Eman Samir Bhaya^{2,3}¹ Department of Mathematics, College of Education for Pure Sciences, University of Babylon, Hillah, Iraq² Department of Mathematics, College of Education, Al-Zahraa University for Women, Karbala, Iraq³ Department of Mathematics, College of Education for Pure Sciences, University of Babylon, Hillah, Iraq

ABSTRACT

The modulus of smoothness is essential for modern analysis and its applications. Is a versatile tool in approximation theory that helps in understanding the properties of approximation methods, characterizing function spaces, and analyzing the convergence and accuracy of numerical algorithms. It helps in determining the optimal number of terms or the optimal choice of basic functions to achieve the desired level of accuracy in approximation of a function and rate of convergence. The smoothness modulus has many applications, including its applications in numerical analysis, particularly in the analysis of numerical methods for solving differential equations, optimization problems, and integral equations. Many papers introduced the ordinary modulus of smoothness with one variable. However, few researchers have tried to approximate functions with multiple variables and mixed modulus. This paper tries to fill in that gap introducing a new $\mathbb{k}$ -mixed modulus of smoothness for measurable functions $f \in L_p([0, 2\pi]^d)$, $0 < p < 1$ with d -variables. It does this by using a new $\mathbb{k}$ -mixed difference and proving some of its approximation properties, like linearity and monotonicity, for the $\mathbb{k}$ -mixed modulus of smoothness of functions belonging to the space L_p using the vector of numbers k_i , $i = 1, \dots, d$. Also, study the approximation of bounded functions with $\mathbb{k}$ -mixed difference and its direct relationship with mixed smoothness.

Keywords: Mixed derivatives, Mixed finite differences, Mixed modulus of smoothness, Monotonicity property, Space of measurable functions

Introduction

Many studies have appeared that dealt with the approximation theory in general, such as,^{1,2} and the modulus of smoothness, in particular, is a powerful tool in approximation theory used to investigate the rate of convergence of approximation methods. The first to develop a concept of dominant mixed modulus of smoothness was Nikol'skii,³ who defined the Sobolev spaces $W_p^\alpha(\Omega)$ defined on an open set $\Omega \subset \mathbb{R}^n$, and it consists of functions that can be p -integrated on Ω up to order α in their partial derivatives. Here, α is a non-negative integer, and $1 < p < \infty$. The space is equipped with a norm that measures the size of the function and its derivatives. The norm of a function $g \in W_p^\alpha(\Omega)$ is given as follows:

$$\|g\|_{W_p^\alpha(\Omega)} = \left(\sum_{|\alpha|} \|D^\alpha g\|_{L_p(\Omega)}^p \right)^{1/p} \quad (1)$$

where α is multi-index, D^α denotes the partial derivative of order $|\alpha|$, and $L_p(\Omega)$ is the p -th Lebesgue space. A dominating mix of smoothness comes from the mixed derivative. Subsequently, studies continued on the dominant mixed smoothness; for example see.⁴⁻⁷ The researchers were also interested in studying the mixed modulus of smoothness it is between mixed metric spaces like.^{8,9}

Received 28 September 2023; revised 1 March 2024; accepted 3 March 2024.
Available online 1 January 2025

* Corresponding author.

E-mail addresses: pure.rehab.amer@uobabylon.edu.iq (R. A. Kamel), emanbhaya@alzahraa.edu.iq (E. S. Bhaya).

<https://doi.org/10.21123/bsj.2024.9726>

2411-7986/© 2025 The Author(s). Published by College of Science for Women, University of Baghdad. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the context of modulus smoothness, precisely that the function g has r -th symmetric difference, according to what follows:

$$\Delta_h^r g(x) = \sum_{j=0}^r \binom{r}{j} (-1)^{r-j} g(x + jh). \quad (2)$$

where, the function $g \in L_p$ has the r -th usual modulus of smoothness which is defined by

$$\omega_r(g, \delta)_p = \sup_{0 < h \leq \delta} \|\Delta_h^r(g, \cdot)\|_p, \delta \geq 0 \quad (3)$$

From Eqs. (2) and (3), it can be developed modulus not in the usual manner, but by defining the modulus of a $\mathbb{k}$ -mixed smoothness of a function $f \in L_p$ as follows:

$$\begin{aligned} \overline{\omega}_{\mathbb{k}}(f, \delta)_p &= \sup_{|h_i| \leq \delta}^d \|\Delta_{h_i}^{\mathbb{k}} f(x_1, x_2, \dots, x_i, \dots, x_d)\|_p, \delta > 0, \text{ or} \\ \underline{\omega}_{\mathbb{k}}(f, \delta)_p &= \sup_{|h_i| \leq \delta}^d \|\underline{\Delta}_{h_i}^{\mathbb{k}} f(x_1, x_2, \dots, x_i, \dots, x_d)\|_p \end{aligned} \quad (4)$$

Then, from Eq. (4), can be defined $\mathbb{k}$ -mixed difference as mentioned below:

$$\begin{aligned} \overline{\Delta}_{h_i}^{\mathbb{k}} f(\mathbb{x}) &= \sup_{i=1}^d \left\{ \Delta_{h_i}^{k_i} f(\mathbb{x}) \right\} \\ &= \sup_{i_1 \leq i \leq d} \left\{ \Delta_{h_{i_1}}^{k_{i_1}} f(\mathbb{x}), \dots, \Delta_{h_{i_i}}^{k_{i_i}} f(\mathbb{x}), \dots, \Delta_{h_{i_d}}^{k_{i_d}} f(\mathbb{x}) \right\}. \end{aligned} \quad (5)$$

Or

$$\begin{aligned} \underline{\Delta}_{h_i}^{\mathbb{k}} f(\mathbb{x}) &= \inf_{i=1}^d \left\{ \Delta_{h_i}^{k_i} f(\mathbb{x}) \right\} \\ &= \inf_{i_1 \leq i \leq d} \left\{ \Delta_{h_{i_1}}^{k_{i_1}} f(\mathbb{x}), \dots, \Delta_{h_{i_i}}^{k_{i_i}} f(\mathbb{x}), \dots, \Delta_{h_{i_d}}^{k_{i_d}} f(\mathbb{x}) \right\}, \end{aligned} \quad (6)$$

where $\mathbb{k} = (k_1, k_2, \dots, k_i, \dots, k_d)$, $h = (h_1, h_2, \dots, h_i, \dots, h_d)$, $\mathbb{x} = (x_1, x_2, \dots, x_i, \dots, x_d)$.

$L_p = L_p([0, 2\pi]^d)$, $0 < p < 1$, be the measurable function's space, and 2π -periodic in all d -variables so that:

$$\|f(\mathbb{x})\|_p = \left(\int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} |f(x_1, x_2, \dots, x_i, \dots, x_d)|^p dx_1 dx_2 \dots dx_i \dots dx_d \right)^{\frac{1}{p}}. \quad (7)$$

Also, $\Delta_{h_i}^{k_i}$ is the standard difference in variable x_i of order k_i with step h_i ; for instance,

$$\begin{aligned} \Delta_{h_i}^1 f(\mathbb{x}) &= \Delta_{h_i}^1 f(x_1, \dots, x_i, \dots, x_d) = f(x_1, \dots, x_i + h_i, \dots, x_d) - f(x_1, \dots, x_i, \dots, x_d), \\ \Delta_{h_i}^{k_i} f(\mathbb{x}) &= \Delta_{h_i}^{k_i} f(\mathbb{x}) \\ &= \sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} f(x_1, \dots, x_i + (k_i - j)h_i, \dots, x_d), \quad k_i \in \mathbb{N}. \end{aligned} \quad (8)$$

Using Eqs. (5) and (6) to obtain

$$\begin{aligned} \Delta_{\gamma h}^{\mathbb{k}} f(\mathbb{x}) &= \sum_{j_{k_1}=0}^{\gamma-1} \dots \sum_{j_{k_i}=0}^{\gamma-1} \dots \sum_{j_{k_d}=0}^{\gamma-1} \sup_{i=1}^d \left\{ \Delta_{h_i}^{k_i} f(x_1, \dots, x_d) \right\} \\ \Delta_{\gamma h}^{\mathbb{k}} f(\mathbb{x}) &= \sum_{j_{k_1}=0}^{\gamma-1} \dots \sum_{j_{k_i}=0}^{\gamma-1} \dots \sum_{j_{k_d}=0}^{\gamma-1} \sup_{i=1}^d \left\{ \Delta_{h_i}^{k_i} f(\mathbb{x}), \dots, \Delta_{h_i}^{k_i} f(\mathbb{x}), \dots, \Delta_{h_d}^{k_d} f(\mathbb{x}) \right\} \end{aligned} \quad (9)$$

Materials and methods

Note: All the following results $\underline{\Delta}_{h_i}^{\mathbb{k}} f(\mathbb{x})$ and $\underline{\omega}_{\mathbb{k}}(f, \delta)_p$ are similar to $\overline{\Delta}_{h_i}^{\mathbb{k}} f(\mathbb{x})$ and $\overline{\omega}_{\mathbb{k}}(f, \delta)_p$ in terms of proof.

Lemma 1:¹⁰ Suppose that $\langle x_l \rangle$ be any sequence; then

$$\left| \sum_l x_l \right|^p \leq 2^{\frac{1}{p}} \sum_l |x_l|^p \quad \text{for } 1 \leq p < \infty;$$

$$\left| \sum_l x_l \right|^p \leq \sum_l |x_l|^p \quad \text{for } 0 < p < 1.$$

Lemma 2:¹¹ For $0 < p < 1$, f and g functions $\in L_p([0, 2\pi]^d)$, then

$$\|(f+g)(x)\|_p \leq (2)^{\frac{1}{p}-1} \|f(x)\|_p + (2)^{\frac{1}{p}-1} \|g(x)\|_p.$$

Lemma 3: Let f and $g \in L_p([0, 2\pi]^d)$, $0 < p < 1$. Then

$$(a) \quad \overline{\Delta_h^k}(f+g)(x) = \overline{\Delta_h^k}f(x) + \overline{\Delta_h^k}g(x);$$

$$(b) \quad \overline{\Delta_h^k}(\overline{\Delta_h^r}f(x)) = \overline{\Delta_h^{k+r}}f(x);$$

$$(c) \quad \left\| \overline{\Delta_h^k}f(x) \right\|_p \leq C(p, k_1) \|f(x)\|_p;$$

$$(d) \quad \lim_{h \rightarrow 0^+} \left\| \overline{\Delta_h^k}f(x) \right\|_p = \lim_{h_1 \rightarrow 0^+} \left\| \Delta_{h_1}^{k_1}f(x) \right\|_p = 0.$$

Proof of:

(a)

$$\begin{aligned} \overline{\Delta_h^k}(f+g)(x) &= \sup_{i=1}^d \left\{ \Delta_{h_i}^{k_i}(f+g)(x) \right\} \\ &= \sup_{i=1}^d \left\{ \Delta_{h_1}^{k_1}(f+g)(x), \Delta_{h_2}^{k_2}(f+g)(x), \dots, \Delta_{h_i}^{k_i}(f+g)(x), \dots, \Delta_{h_d}^{k_d}(f+g)(x) \right\} \\ &= \sup_{i=1}^d \left\{ \begin{aligned} &\sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} (f+g)(x_1 + (k_1-j)h_1, x_2, \dots, x_i, \dots, x_d), \\ &\sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} (f+g)(x_1, x_2 + (k_2-j)h_2, \dots, x_i, \dots, x_d), \dots, \\ &\sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} (f+g)(x_1, x_2, \dots, x_i + (k_i-j)h_i, \dots, x_d), \dots, \\ &\sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} (f+g)(x_1, x_2, \dots, x_i, \dots, x_d + (k_d-j)h_d) \end{aligned} \right\} \end{aligned}$$

From the properties of finite difference, the following can be concluded

$$\overline{\Delta_h^k}(f+g)(x) = \sup_{i=1}^d \left\{ \begin{aligned} &\sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} f(x_1 + (k_1-j)h_1, x_2, \dots, x_i, \dots, x_d) + \\ &\sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} g(x_1 + (k_1-j)h_1, x_2, \dots, x_i, \dots, x_d), \\ &\sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} f(x_1, x_2 + (k_2-j)h_2, \dots, x_i, \dots, x_d) + \\ &\sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} g(x_1, x_2 + (k_2-j)h_2, \dots, x_i, \dots, x_d), \dots, \\ &\sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} f(x_1, x_2, \dots, x_i + (k_i-j)h_i, \dots, x_d) + \\ &\sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} g(x_1, x_2, \dots, x_i + (k_i-j)h_i, \dots, x_d), \dots, \\ &\sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} f(x_1, x_2, \dots, x_i, \dots, x_d + (k_d-j)h_d) + \\ &\sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} g(x_1, x_2, \dots, x_i, \dots, x_d + (k_d-j)h_d) \end{aligned} \right\}$$

Then from properties of supremum,

$$\begin{aligned} \overline{\Delta_h^k} (f+g)(x) &= \sup_{i=1}^d \left\{ \begin{aligned} &\sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} f(x_1 + (k_1 - j)h_1, x_2, \dots, x_i, \dots, x_d), \\ &\sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} f(x_1, x_2 + (k_2 - j)h_2, \dots, x_i, \dots, x_d), \dots, \\ &\sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} f(x_1, x_2, \dots, x_i + (k_i - j)h_i, \dots, x_d), \dots, \\ &\sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} f(x_1, x_2, \dots, x_i, \dots, x_d + (k_d - j)h_d) \end{aligned} \right\} \\ &\quad + \sup_{i=1}^d \left\{ \begin{aligned} &\sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} g(x_1 + (k_1 - j)h_1, x_2, \dots, x_i, \dots, x_d), \\ &\sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} g(x_1, x_2 + (k_2 - j)h_2, \dots, x_i, \dots, x_d), \dots, \\ &\sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} g(x_1, x_2, \dots, x_i + (k_i - j)h_i, \dots, x_d), \dots, \\ &\sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} g(x_1, x_2, \dots, x_i, \dots, x_d + (k_d - j)h_d) \end{aligned} \right\} \\ &= \overline{\Delta_h^k} f(x) + \overline{\Delta_h^k} g(x). \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \overline{\Delta_h^k} (\overline{\Delta_h^r} f(x)) &= \overline{\Delta_h^k} \left(\sup_{i=1}^d \left\{ \Delta_{h_i}^{r_1} f(x) \right\} \right) \\ &= \overline{\Delta_h^k} \left(\sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \right) \\ &= \sup_{i=1}^d \left\{ \Delta_{h_i}^{k_i} \left(\sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \right) \right\} \\ \overline{\Delta_h^k} (\overline{\Delta_h^r} f(x)) &= \sup_{i=1}^d \left\{ \begin{aligned} &\Delta_{h_1}^{k_1} \left(\sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \right), \\ &\Delta_{h_2}^{k_2} \left(\sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \right), \\ &\dots, \\ &\Delta_{h_i}^{k_i} \left(\sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \right), \\ &\dots, \\ &\Delta_{h_d}^{k_d} \left(\sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \right) \end{aligned} \right\} \end{aligned}$$

Choose $r_1, r_2, \dots, r_i, \dots, r_d$ s.t. $\Delta_{h_1}^{r_1} f(x) = \sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\}$

$$\begin{aligned} \Delta_{h_2}^{r_2} f(x) &= \sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \\ &\vdots \\ \Delta_{h_i}^{r_i} f(x) &= \sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \\ &\vdots \\ \Delta_{h_d}^{r_d} f(x) &= \sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \end{aligned}$$

Then

$$\overline{\Delta_h^k} (\overline{\Delta_h^r} f(x)) = \sup_{i=1}^d \left\{ \Delta_{h_1}^{k_1} \left(\Delta_{h_1}^{r_1} f(x) \right), \Delta_{h_2}^{k_2} \left(\Delta_{h_2}^{r_2} f(x) \right), \dots, \Delta_{h_i}^{k_i} \left(\Delta_{h_i}^{r_i} f(x) \right), \dots, \Delta_{h_d}^{k_d} \left(\Delta_{h_d}^{r_d} f(x) \right) \right\}$$

From properties of finite difference, the following result

$$\begin{aligned}
 \overline{\Delta_h^k}(\overline{\Delta_h^r} f(x)) &= \sup_{i=1}^d \left\{ \Delta_{h_1}^{k_1+r_1} f(x), \Delta_{h_2}^{k_2+r_2} f(x), \dots, \Delta_{h_i}^{k_i+r_i} f(x), \dots, \Delta_{h_1}^{k_1+r_1} f(x), \Delta_{h_d}^{k_d+r_d} f(x) \right\} \\
 &= \overline{\Delta_h^{k+r}} f(x). \\
 \text{(c) } \left\| \overline{\Delta_h^k} f(x) \right\|_p &= \left\| \sup_{i=1}^d \left\{ \Delta_{h_i}^{k_i} f(x) \right\} \right\|_p \\
 &= \left\| \sup_{i=1}^d \left\{ \Delta_{h_1}^{r_1} f(x), \Delta_{h_2}^{r_2} f(x), \dots, \Delta_{h_i}^{r_i} f(x), \dots, \Delta_{h_d}^{r_d} f(x) \right\} \right\|_p \\
 &= \left\| \sup_{i=1}^d \left\{ \sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} f(x_1 + (k_1 - j)h_1, x_2, \dots, x_i, \dots, x_d), \right. \right. \\
 &\quad \left. \sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} f(x_1, x_2 + (k_2 - j)h_2, \dots, x_i, \dots, x_d), \dots, \right. \\
 &\quad \left. \sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} f(x_1, x_2, \dots, x_i + (k_i - j)h_i, \dots, x_d), \dots, \right. \\
 &\quad \left. \sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} f(x_1, x_2, \dots, x_i, \dots, x_d + (k_d - j)h_d) \right\} \Bigg\|_p \\
 \left\| \overline{\Delta_h^k} f(x) \right\|_p &= \left(\int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \left| \sup_{i=1}^d \left\{ \sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} f(x_1 + (k_1 - j)h_1, x_2, \dots, x_i, \dots, x_d), \right. \right. \right. \\
 &\quad \left. \sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} f(x_1, x_2 + (k_2 - j)h_2, \dots, x_i, \dots, x_d), \dots, \right. \\
 &\quad \left. \sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} f(x_1, x_2, \dots, x_i + (k_i - j)h_i, \dots, x_d), \dots, \right. \\
 &\quad \left. \sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} f(x_1, x_2, \dots, x_i, \dots, x_d + (k_d - j)h_d) \right\} \Bigg|^p dx_1 \dots dx_i \dots dx_d \Bigg)^{\frac{1}{p}} \\
 \text{Choose } k_i \text{ s.t. } \overline{\Delta_{h_i}^{k_i}} f(x) &= \sup_{i=1}^d \left\{ \sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} f(x_1 + (k_1 - j)h_1, x_2, \dots, x_i, \dots, x_d), \right. \\
 &\quad \left. \sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} f(x_1, x_2 + (k_2 - j)h_2, \dots, x_i, \dots, x_d), \dots, \right. \\
 &\quad \left. \sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} f(x_1, x_2, \dots, x_i + (k_i - j)h_i, \dots, x_d), \dots, \right. \\
 &\quad \left. \sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} f(x_1, x_2, \dots, x_i, \dots, x_d + (k_d - j)h_d) \right\}
 \end{aligned}$$

From Lemma 1

$$\begin{aligned}
 \left\| \overline{\Delta_h^k} f(x) \right\|_p &\leq \left(\int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} \sum_{j=0}^{k_1} \left| \binom{k_1}{j} \right|^p |f(x_1, x_2, \dots, x_1 + (k_1 - j)h_1, \dots, x_d)|^p dx_1 \dots dx_i \dots dx_d \right)^{\frac{1}{p}} \\
 &= \left(\sum_{j=0}^{k_1} \left| \binom{k_1}{j} \right|^p \int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} |f(x_1, x_2, \dots, x_1 + (k_1 - j)h_1, \dots, x_d)|^p dx_1 \dots dx_i \dots dx_d \right)^{\frac{1}{p}} \\
 &\leq C(p, k_1) \left(\int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} |f(x_1, x_2, \dots, x_1 + (k_1 - j)h_1, \dots, x_d)|^p dx_1 \dots dx_i \dots dx_d \right)^{\frac{1}{p}} \\
 &\leq C(p, k_1) \|f(x)\|_p.
 \end{aligned}$$

$$\begin{aligned}
 \text{(d) } \lim_{h \rightarrow 0^+} \left\| \overline{\Delta_h^k} f(x) \right\|_p &= \lim_{h \rightarrow 0^+} \left\| \sup_{i=1}^d \left(\Delta_{h_i}^{k_i} f(x) \right) \right\|_p \\
 &= \lim_{h \rightarrow 0^+} \left\| \sup_{i=1}^d \left(\Delta_{h_1}^{k_1} f(x), \Delta_{h_2}^{k_2} f(x), \dots, \Delta_{h_i}^{k_i} f(x), \dots, \Delta_{h_d}^{k_d} f(x) \right) \right\|_p,
 \end{aligned}$$

where $k_1 \in \mathbb{N}$ such that

$$\begin{aligned}\Delta_{h_1}^{k_1} f(x) &= \sup_{i=1}^d \left\{ \Delta_{h_1}^{k_1} f(x), \Delta_{h_2}^{k_2} f(x), \dots, \Delta_{h_i}^{k_i} f(x), \dots, \Delta_{h_d}^{k_d} f(x) \right\} \\ &= \lim_{h_1 \rightarrow 0^+} \left\| \Delta_{h_1}^{k_1} f(x) \right\|_p \\ &= \lim_{h_1 \rightarrow 0^+} \left\| \sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} f(x_1, x_2, \dots, x_1 + (k_1 - j)h_1, \dots, x_d) \right\|_p \\ \Delta_{h_1}^{k_1} f(x) &= \lim_{h_1 \rightarrow 0^+} \left\| \left((-1)^0 \frac{k_1!}{0!(k_1-0)!} f(x_1, x_2, \dots, x_1 + k_1 h_1, \dots, x_d) + (-1)^1 \frac{k_1!}{1!(k_1-1)!} f(x_1, x_2, \dots, x_1 \right. \right. \\ &\quad \left. \left. + (k_1 - 1)h_1, \dots, x_d) + \dots + (-1)^j \frac{k_1!}{j!(k_1-j)!} f(x_1, x_2, \dots, x_1 + (k_1 - j)h_1, \dots, x_d) \right. \right. \\ &\quad \left. \left. + \dots + (-1)^{k_1} \frac{k_1!}{k_1!(k_1-k_1)!} f(x_1, x_2, \dots, x_1 + (k_1 - k_1)h_1, \dots, x_d) \right) \right\|_p \\ &= \lim_{k_1 \rightarrow \infty} \left\| \sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} f(x_1, x_2, \dots, x_1, \dots, x_d) \right\|_p = 0\end{aligned}$$

Results and discussion

In the following result, can be summarized the major characteristics of modulus of a k -mixed smoothness of $f \in L_p([0, 2\pi]^d)$, $0 < p < 1$

Theorem 1: Let f and $g \in L_p([0, 2\pi]^d)$, $0 < p < 1$, and $k = (k_1, k_2, \dots, k_i, \dots, k_d)$, $k_i \in \mathbb{N}$. Then

- (a) $\lim_{\delta \rightarrow 0^+} \overline{\omega}_k(f, \delta)_p = 0$;
- (b) $\overline{\omega}_k(f + g, \delta)_p \leq 2^{\frac{1}{p}-1} (\overline{\omega}_k(f, \delta)_p + \overline{\omega}_k(g, \delta)_p)$;
- (c) $\overline{\omega}_k(f, \delta)_p \leq \overline{\omega}_k(f, t)_p$, for $0 < \delta < t$;
- (d) $\frac{\overline{\omega}_k(f, \delta)_p}{\delta^k} \leq \frac{\overline{\omega}_k(f, t)_p}{t^k}$, for $0 < t \leq \delta \leq 1$;
- (e) $\overline{\omega}_k(f, \gamma \delta)_p \leq \gamma^{k_1} \overline{\omega}_k(f, \delta)_p$, for $\gamma > 1$ and $\gamma \in \mathbb{N}$, where $k_1 = \max |k_i|$, $i = 1, \dots, d$

Proof of:

(a)

$$\begin{aligned}\overline{\omega}_k(f, \delta)_p &= \sup_{|h_i| \leq \delta} \left\| \Delta_h^k f(x_1, x_2, \dots, x_i, \dots, x_d) \right\|_p \\ \lim_{\delta \rightarrow 0^+} \overline{\omega}_k(f, \delta)_p &= \lim_{\delta \rightarrow 0^+} \sup_{|h_i| \leq \delta} \left\| \Delta_h^k f(x_1, x_2, \dots, x_i, \dots, x_d) \right\|_p\end{aligned}$$

From Lemma 3(d)

$$\lim_{\delta \rightarrow 0^+} \overline{\omega}_k(f, \delta)_p = 0.$$

(b)

$$\begin{aligned}\overline{\omega}_k(f + g, \delta)_p &= \sup_{|h_i| \leq \delta} \left\| \Delta_h^k (f + g)(x) \right\|_p \\ &= \sup_{|h_i| \leq \delta} \left\| \Delta_h^k (f + g)(x_1, \dots, x_d) \right\|_p \\ &= \sup_{|h_i| \leq \delta} \left\| \sup_{i=1}^d \left(\Delta_{h_i}^{k_i} (f + g) \right)(x_1, \dots, x_d) \right\|_p \\ &= \sup_{|h_i| \leq \delta} \left\| \left(\int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} \left| \sup_{i=1}^d \left\{ \Delta_{h_1}^{k_1} (f + g)(x_1, \dots, x_d), \Delta_{h_2}^{k_2} (f + g)(x_1, \dots, x_d), \right. \right. \right. \right. \\ &\quad \left. \left. \left. \dots, \Delta_{h_i}^{k_i} (f + g)(x_1, \dots, x_d), \dots, \Delta_{h_d}^{k_d} (f + g)(x_1, \dots, x_d) \right\} \right|^p dx_1 \dots dx_d \right)^{\frac{1}{p}} \right\|_p\end{aligned}$$

$$\bar{\omega}_k(f+g, \delta)_p = \sup_{|h_i| \leq \delta} \left(\int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} \sup_{i=1}^d \left| \sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} (f+g)(x_1 + (k_1-j)h_1, x_2, \dots, x_i, \dots, x_d), \right. \right. \\ \left. \left. \sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} (f+g)(x_1, x_2 + (k_2-j)h_2, \dots, x_i, \dots, x_d), \dots, \right. \right. \\ \left. \left. \sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} (f+g)(x_1, x_2, \dots, x_i + (k_i-j)h_i, \dots, x_d), \dots, \right. \right. \\ \left. \left. \sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} (f+g)(x_1, x_2, \dots, x_i, \dots, x_d + (k_d-j)h_d) \right| dx_1 \dots dx_d \right)^{\frac{1}{p}}$$

Can be obtained from [Lemmas 2](#) and [3\(a\)](#),

$$\bar{\omega}_k(f+g, \delta)_p \leq 2^{\frac{1}{p}-1} \sup_{|h_i| \leq \delta} \left(\int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} \sup_{i=1}^d \left| \sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} f(x_1 + (k_1-j)h_1, x_2, \dots, x_i, \dots, x_d), \right. \right. \\ \left. \left. \sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} f(x_1, x_2 + (k_2-j)h_2, \dots, x_i, \dots, x_d), \dots, \right. \right. \\ \left. \left. \sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} f(x_1, x_2, \dots, x_i + (k_i-j)h_i, \dots, x_d), \dots, \right. \right. \\ \left. \left. \sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} f(x_1, x_2, \dots, x_i, \dots, x_d + (k_d-j)h_d) \right| dx_1 \dots dx_i \dots dx_d \right)^{\frac{1}{p}} \\ + \left(\int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} \sup_{i=1}^d \left| \sum_{j=0}^{k_1} (-1)^j \binom{k_1}{j} g(x_1 + (k_1-j)h_1, x_2, \dots, x_i, \dots, x_d), \right. \right. \\ \left. \left. \sum_{j=0}^{k_2} (-1)^j \binom{k_2}{j} g(x_1, x_2 + (k_2-j)h_2, \dots, x_i, \dots, x_d), \dots, \right. \right. \\ \left. \left. \sum_{j=0}^{k_i} (-1)^j \binom{k_i}{j} g(x_1, x_2, \dots, x_i + (k_i-j)h_i, \dots, x_d), \dots, \right. \right. \\ \left. \left. \sum_{j=0}^{k_d} (-1)^j \binom{k_d}{j} g(x_1, x_2, \dots, x_i, \dots, x_d + (k_d-j)h_d) \right| dx_1 \dots dx_i \dots dx_d \right)^{\frac{1}{p}}$$

$$\bar{\omega}_k(f+g, \delta)_p \leq 2^{\frac{1}{p}-1} \left(\sup_{|h_i| \leq \delta} \left\| \Delta_h^k f(x_1, x_2, \dots, x_i, \dots, x_d) \right\|_p + \sup_{|h_i| \leq \delta} \left\| \Delta_h^k g(x_1, x_2, \dots, x_i, \dots, x_d) \right\|_p \right)$$

$$\bar{\omega}_k(f+g, \delta)_p \leq 2^{\frac{1}{p}-1} \left(\sup_{|h_i| \leq \delta} \left\| \Delta_h^k f(x_1, \dots, x_d) \right\|_p + \sup_{|h_i| \leq \delta} \left\| \Delta_h^k g(x_1, \dots, x_d) \right\|_p \right)$$

$$\leq 2^{\frac{1}{p}-1} (\bar{\omega}_k(f, \delta)_p + \bar{\omega}_k(g, \delta)_p).$$

(c)

$$\bar{\omega}_k(f, \delta)_p = \sup_{|h_i| \leq \delta} \sup_{i=1}^d \left\| \overline{\Delta_{h_i}^k} f(x) \right\|_p$$

$$\bar{\omega}_k(f, \delta)_p = \sup_{|h_i| \leq \delta} \sup_{i=1}^d \left\| \overline{\Delta_{h_i}^k} f(x_1, x_2, \dots, x_i, \dots, x_d) \right\|_p$$

since $0 < \delta < t$, then

$$\bar{\omega}_k(f, \delta)_p \leq \sup_{|h_i| \leq t} \sup_{i=1}^d \left\| \overline{\Delta_{h_i}^k} f(x_1, x_2, \dots, x_i, \dots, x_d) \right\|_p$$

$$= \bar{\omega}_k(f, t)_p.$$

(d) Suppose that $k = \max |k_i|_{1 \leq i \leq d}$

Since $\delta \leq 1$, then $\frac{1}{\delta^k} \leq \frac{1}{t^k}$.

From (c) $\rightarrow \frac{\bar{\omega}_k(f, \delta)_p}{\delta^k} \leq \frac{\bar{\omega}_k(f, t)_p}{t^k}$, for $0 < \delta < t$.

(e) By applying Eq. (9), to obtain

$$\Delta_{\gamma h}^k f(x) = \sum_{j_{k_1}=0}^{\gamma-1} \dots \sum_{j_{k_m}=0}^{\gamma-1} \dots \sum_{j_{k_d}=0}^{\gamma-1} \sup_{i=1}^d \left\{ \Delta_{h_1}^{k_1} f(x), \dots, \Delta_{h_i}^{k_i} f(x), \dots, \Delta_{h_d}^{k_d} f(x) \right\},$$

where $k_i = \max |k_i|$, $i = 1, \dots, d$. k_i is one component of $(k_1, \dots, k_d)$, then

$$\Delta_{\gamma h_1}^{k_1} f(x) = \sum_{j_{k_1}=0}^{\gamma-1} \dots \sum_{j_{k_m}=0}^{\gamma-1} \dots \sum_{j_{k_d}=0}^{\gamma-1} \Delta_{h_1}^{k_1} f(x_1 \dots x_1 \dots x_d)$$

$$\Delta_{\gamma h_1}^{k_1} f(x) = \sum_{j_{k_1}=0}^{\gamma-1} \dots \sum_{j_{k_m}=0}^{\gamma-1} \dots \sum_{j_{k_1}=0}^{\gamma-1} \sum_{j=0}^{k_1} (-1)^j \left(\frac{k_1}{j} \right) f(x_1, \dots, x_1 + (k_1 - j) h_1 + (k_1 - j_{k_1}) h_1$$

$$+ \dots + (k_1 - j_{k_1}) h_1, \dots, x_d)$$
(10)

By induction on k_1 can be proved the above definition. If $k_1 = 1 \rightarrow$

$$\Delta_{\gamma h_1}^1 f(x) = \Delta_{\gamma h_1}^1 f(x_1, \dots, x_1, \dots, x_d) = f(x_1, \dots, x_1 + \gamma h_1, \dots, x_d) - f(x_1, \dots, x_1, \dots, x_d)$$

$$= \sum_{j=0}^{\gamma-1} [f(x_1, \dots, x_1 + j h_1 + h_1, \dots, x_d) - f(x_1, \dots, x_1 + j h_1, \dots, x_d)]$$

$$\Delta_{\gamma h_1}^1 f(x) = \sum_{j=0}^{\gamma-1} \Delta_{h_1} f(x_1, \dots, x_1 + j h_1, \dots, x_d)$$

For arbitrary $\mathbf{k}_l \in \mathbb{N}$ and by Lemma 3(b)→

$$\begin{aligned}
 \Delta_{\gamma h_l}^{\mathbf{k}_l+1} f(\mathbf{x}) &= \Delta_{\gamma h_l}^{\mathbf{k}_l} \left[\Delta_{\gamma h_l}^1 f(\mathbf{x}) \right] \\
 &= \sum_{j_{k_1}=0}^{\gamma-1} \cdots \sum_{j_{k_m}=0}^{\gamma-1} \cdots \sum_{j_{k_l}=0}^{\gamma-1} \Delta_{h_l}^{\mathbf{k}_l} \left[\Delta_{\gamma h_l}^1 (f(x_1, \dots, x_l + (\mathbf{k}_l - j_{k_1}) h_l + (\mathbf{k}_l - j_{k_2}) h_l + \cdots + (\mathbf{k}_l - j_{k_l}) h_l, \dots, x_d)) \right] \\
 &= \sum_{j_{k_1}=0}^{\gamma-1} \cdots \sum_{j_{k_m}=0}^{\gamma-1} \cdots \sum_{j_{k_l}=0}^{\gamma-1} \Delta_{h_l}^{\mathbf{k}_l} \left[\sum_{j_{k_l+1}=0}^{\gamma-1} \Delta_{h_l} (f(x_1, \dots, x_l + (\mathbf{k}_l - j_{k_1}) h_l + (\mathbf{k}_l - j_{k_2}) h_l \right. \\
 &\quad \left. + \cdots + (\mathbf{k}_l - j_{k_l}) h_l + (\mathbf{k}_l - j_{k_l+1}) h_l, \dots, x_d)) \right] \\
 &= \sum_{j_{k_1}=0}^{\gamma-1} \cdots \sum_{j_{k_m}=0}^{\gamma-1} \cdots \sum_{j_{k_l}=0}^{\gamma-1} \sum_{j_{k_l+1}=0}^{\gamma-1} \Delta_{h_l}^{\mathbf{k}_l+1} f(x_1, \dots, x_l + (\mathbf{k}_l - j_{k_1}) h_l + (\mathbf{k}_l - j_{k_2}) h_l \\
 &\quad + \cdots + (\mathbf{k}_l - j_{k_l}) h_l + (\mathbf{k}_l - j_{k_l+1}) h_l, \dots, x_d)
 \end{aligned}$$

Form Eq. (10) → for $h_l, |h_l| \leq \delta$

$$\begin{aligned}
 \left| \Delta_{\gamma h_l}^{\mathbf{k}_l} f(\mathbf{x}) \right| &\leq \sum_{j_{k_2}=0}^{\gamma-1} \cdots \sum_{j_{k_m}=0}^{\gamma-1} \cdots \sum_{j_{k_l}=0}^{\gamma-1} \left\| \Delta_{h_l}^{\mathbf{k}_l} f(x_1, \dots, x_l + (\mathbf{k}_l - j_{k_2}) h_l + (\mathbf{k}_l - j_{k_3}) h_l + \cdots + (\mathbf{k}_l - j_{k_l}) h_l, \dots, x_d) \right\|_p \\
 &\leq \gamma^{\mathbf{k}_l} \omega_{\mathbf{k}_l}(f, \delta)_p.
 \end{aligned}$$

Conclusion

This paper studies a new mixed difference for $\mathbb{k}$ -vectors in d -variables. Thus, a new modulus of $\mathbb{k}$ -mixed smoothness is obtained, and the fundamental modulus characteristics of $\mathbb{k}$ -mixed smoothness are also proved. Also, it can be concluded that the $\mathbb{k}$ -mixed modulus of smoothness achieves most of the inequalities achieved by the usual modulus of smoothness, such as monotonicity and linearity, in addition to a basic inequality, which is the boundness of functions belonging to the space L_p , $0 < p < 1$, with a mixed difference of order $\mathbb{k}$.

Acknowledgment

Thank God and everyone who supported us to accomplish this work.

Authors' declaration

- Conflicts of Interest: None.
- No animal studies are present in the manuscript.
- No human studies are present in the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee at University of Babylon.

Authors' contribution statement

This work was carried out in collaboration between all authors. E. S. B, wrote and edited the manuscript with revisions idea. R. A. K, analysis the data with manuscript design. All authors read and approved the final manuscript.

References

1. Kasim NM. New Theorems in approximation theory. Baghdad Sci J. 2010;7(2):1056–60. <https://doi.org/10.21123/bsj.2010.7.2.1056-1060>.

2. Aal-Rkhais HA, Kamil AH, Al Oweidi KF. The approximation of weighted Hölder functions by Fourier-Jacobi polynomials to the singular Sturm-Liouville operator. *Baghdad Sci J.* 2022;19(6):1387–92. <https://dx.doi.org/10.21123/bsj.2022.6128>.
3. Nikolsky SM. Functions with dominating mixed derivatives satisfying multiple holder conditions. *Sib Math J.* 1963;4(6):1342–64.
4. Najafov AM, Gasimova AM. On some differential properties of functions in Lizorkin-Triebel-Morrey spaces with dominant mixed derivatives. *Adv Appl Math Sci.* 2021;20(9):1909–21.
5. Akgün R. Fractional order mixed difference operator and its applications in angular approximation. *Hacet J Math Stat.* 2020;49(5):1594–610. <https://doi.org/10.15672/hujms.569410>.
6. Zhao J, Du W-S, Chen Y. New generalizations and results in shift-invariant subspaces of mixed-norm Lebesgue spaces $L_p(\mathbb{R}^d)$. *Mathematics.* 2021;9(3):1–10. <https://doi.org/10.3390/math9030227>.
7. Zhao J, Kostic M, Du W-S. On new decomposition theorems for mixed-norm besove spaces with ingredient modulus of smoothness. *Symmetry.* 2023;15(3):1–17. <https://doi.org/10.3390/sym15030642>.
8. Potapov MK, Simonov BV. Estimates of partial moduli of smoothness in metrics of $L_{p_1\infty}$ and $L_{\infty p_2}$ through partial moduli of smoothness in metrics of $L_{p_1 p_2}$. *Moscow Univ Math Bull.* 2020;75:1–15. <https://doi.org/10.3103/S0027132220010015>.
9. Potapov MK, Simonov BV. Refinement of relations between mixed moduli of smoothness in the metrics of L_p and L_∞ . *Math Notes.* 2021;110:368–85. <https://doi.org/10.1134/S0001434621090054>.
10. Zygmund A. Trigonometric series. 3rd Ed. Volume I & II. United Kingdom: Cambridge University Press; 1958;728p. <https://doi.org/10.1017/CBO9781316036587>.
11. Castillo RE, Rafeiro H. An introductory course in Lebesgue spaces. 1st edition. Switzerland: Springer; 2016. 461p. <https://doi.org/10.1007/978-3-319-30034-4>.

مقياس النعومة $\mathbb{K}$ - المختلط

رحاب عامر كامل¹، ايمان سمير بهيه^{2,3}

¹ قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعة بابل، الحلة، العراق.

² قسم الرياضيات، كلية التربية، جامعة الزهراء للنبات، كربلاء، العراق.

³ قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعة بابل، الحلة، العراق.

الخلاصة

يعتبر معامل او مقياس النعومة من المواضيع الاساسية والمهمة في التحليل الرياضي وتطبيقاته المتعددة. حيث يستخدم قياس نعومة الدوال معامل النعومة لاستنتاج الخصائص التقريبية للدوال التي يتعذر ايجاد قيمة دقيقة لها. لمقياس النعومة دور اساسي في العديد من مجالات الرياضيات وله تطبيقات مختلفة في نظرية التقريب و في التحليل العددي وخاصة في تحليل الطرق العددية لحل المعادلات التفاضلية ومشاكل التحسين والمعادلات التكاملية و معدل التقارب. تناول العديد من الباحثين في دراستهم معامل النعومة لدالة ذات متغير واحد ولكن قلة من الباحثين الذين تناولوا مقياس النعومة لدالة ذات عدة متغيرات. يهدف هذا البحث الى ملئ هذه الفجوة حيث عرف مقياس نعومة مختلط جديد لدالة تنتمي لفضاء الدوال القابلة للتكامل لبيكيا ذات d من المتغيرات. كما اثبتت العديد من الخصائص الاساسية لمقياس النعومة ولكن باستخدام مقياس نعومة مختلط لدالة ذات فرق منتهي مختلط و d من المتغيرات كالخاصية الخطية والرتابة اضافة الى ذلك دراسة خاصية اساسية وهي المحدودية لدوال تنتمي لفضاء الدوال القابلة للتكامل والقياس.

الكلمات المفتاحية: المشتقات المختلطة، الفرق المنتهي المختلط، معامل النعومة المختلط، الخاصية الرتيبة، فضاء الدوال القابلة للقياس.