

2025

Outcome for the Case (11,10; 1)

Waleed Khalid Jaber

Department of Mathematics, College of Education for Pure Science Ibn-Al-Haitham, University of Baghdad, Baghdad, Iraq AND Department of Mathematics, College of Computer Sciences and Mathematics, University of Thi-Qar, Nassiriyah, Iraq, waleedkhalidj@utq.edu.iq

Niran Sabah Jasim

Department of Mathematics, College of Education for Pure Science Ibn-Al-Haitham, University of Baghdad, Baghdad, Iraq

Follow this and additional works at: <https://bsj.researchcommons.org/home>

How to Cite this Article

Jaber, Waleed Khalid and Jasim, Niran Sabah (2025) "Outcome for the Case (11,10; 1)," *Baghdad Science Journal*: Vol. 22: Iss. 2, Article 25.

DOI: <https://doi.org/10.21123/bsj.2024.9905>

This Article is brought to you for free and open access by Baghdad Science Journal. It has been accepted for inclusion in Baghdad Science Journal by an authorized editor of Baghdad Science Journal.



RESEARCH ARTICLE

Outcome for the Case (11,10; 1)

Waleed Khalid Jaber^{1,2,*}, Niran Sabah Jasim¹¹ Department of Mathematics, College of Education for Pure Science Ibn-Al-Haitham, University of Baghdad, Baghdad, Iraq² Department of Mathematics, College of Computer Sciences and Mathematics, University of Thi-Qar, Nassiriyah, Iraq

ABSTRACT

Through this work R is a commutative ring with 1, $\mathcal{F}$ is a free R -module and $\mathcal{D}_i$ is the divided power algebra of degree i . A partition of length $\ell(\lambda) = n$ is a sequence $(\lambda) = (\lambda_1, \lambda_2, \dots, \lambda_n)$ of non-negative integers in non-increasing order $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > 0$. The weight of a partition λ is $|\lambda| = \lambda_1 + \lambda_2 + \dots + \lambda_n$. A relative sequence is a pair (λ, μ) of sequences such that $\mu \leq \lambda$ and denoted by λ/μ . If both (λ) and μ are partitions then the relative sequence (λ/μ) is called a skew partition. Let Z_{21} be the free generator of divided power algebra $\mathcal{D}(Z_{21})$ in one generator. The divided power element $Z_{21}^{(e)}$ of degree e of the free generator Z_{21} acts on $\mathcal{D}_{p+e} + \mathcal{D}_{q-e}$ by place polarization of degree e from place 1 to place 2. The graded algebra $\mathcal{A} = \mathcal{D}(Z_{21})$ acts on the graded module $\mathcal{M} = \mathcal{D}_{p+e} + \mathcal{D}_{q-e} = \sum \mathcal{M}_{q-e}$, the degree of the second factor determines the grading. $\mathcal{M}$ is a graded left $\mathcal{A}$ -module, where for $w = Z_{21}^{(e)} \in \mathcal{A}$ and $v \in \mathcal{D}_{p+e} + \mathcal{D}_{q-e}$, so we have: $w(v) = Z_{21}^{(e)}(v) = \partial_{21}^{(e)}$. This article surveys the exactness of the Weyl resolution of the sequence for the case (11,10; 1) after getting the contracting homotopy of $\{\mathcal{S}_i\}$; where $i = 0, 1, \dots, 8$, and the terms of the characteristic-free resolution for it.

Keywords: Divided power algebra, Letter place, Overlap, Place polarizations, Resolution, Weyl module

Introduction

Writer in¹ discuss $\mathcal{K}_{\lambda/\mu}\mathcal{F}$ as:

$$\lambda/\mu = \begin{array}{c} \begin{array}{|c|} \hline t \\ \hline \end{array} \begin{array}{|c|} \hline \\ \hline \end{array} \begin{array}{|c|} \hline p \\ \hline \end{array} \\ \begin{array}{|c|} \hline \\ \hline \end{array} \begin{array}{|c|} \hline q \\ \hline \end{array} \end{array}$$

For $\mathcal{K}_{\lambda/\mu}\mathcal{F} = \text{Im}(d'_{\lambda/\mu})$ where $d'_{\lambda/\mu}: \mathcal{DF} \rightarrow \bigwedge \mathcal{F}$ (Weyl map), so

$$\sum \mathcal{D}_{p+k} \otimes \mathcal{D}_{q-k} \xrightarrow{\square} \mathcal{D}_p \otimes \mathcal{D}_q \xrightarrow{d'_{\lambda/\mu}} \mathcal{K}_{\lambda/\mu} \rightarrow 0$$

Searchers in² define the representation of the group. For case (11,10;1) the authors treatise this treatise as an application of the Weyl resolution in cases (4,4,3),³ this can be employed for the differential subordination and superordination, **fuzzy maximal sub-modules**, almost projective semimodules and semigroup ideals and right n -derivation in 3-prime near-rings.⁴⁻⁷ Authors in^{8,9} study the **hyper fuzzy AT-ideals of AT-algebra and spectrum of secondary submodules** which can be also apply to this idea.

Main upshots

As in^{1,5} for the case (11,10;1) gainReceived 11 October 2023; revised 4 March 2023; accepted 6 March 2024.
Available online xx xxxx 2025

* Corresponding author.

E-mail addresses: waleedkhalidj@utq.edu.iq (W. K. Jaber), niraan.s.j@ihcoedu.uobaghdad.edu.iq (N. S. Jasim).<https://doi.org/10.21123/bsj.2024.9905>

2411-7986/© 2025 The Author(s). Published by College of Science for Women, University of Baghdad. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

$$\mathcal{M}_0 = \mathcal{D}_{10} \otimes \mathcal{D}_{10}$$

$$\mathcal{M}_1 = \mathcal{Z}_{21}^{(b)} x \mathcal{D}_{10+b} \otimes \mathcal{D}_{10-b}; \text{ where } 2 \leq b \leq 10$$

$$\mathcal{M}_2 = \mathcal{Z}_{21}^{(b_1)} x \mathcal{Z}_{21}^{(b_2)} x \mathcal{D}_{10+|b|} \otimes \mathcal{D}_{10-|b|}; \text{ where } 3 \leq |b| = b_1 + b_2 \leq 10 \text{ and } b_1 \geq 2$$

$$\mathcal{M}_3 = \mathcal{Z}_{21}^{(b_1)} x \mathcal{Z}_{21}^{(b_2)} x \mathcal{Z}_{21}^{(b_3)} x \mathcal{D}_{10+|b|} \otimes \mathcal{D}_{10-|b|}; \text{ where } 4 \leq |b| = \sum_{i=1}^3 b_i \leq 10 \text{ and } b_1 \geq 2$$

$$\mathcal{M}_4 = \mathcal{Z}_{21}^{(b_1)} x \mathcal{Z}_{21}^{(b_2)} x \mathcal{Z}_{21}^{(b_3)} x \mathcal{Z}_{21}^{(b_4)} x \mathcal{D}_{10+|b|} \otimes \mathcal{D}_{10-|b|}; \text{ where } 5 \leq |b| = \sum_{i=1}^4 b_i \leq 10 \text{ and } b_1 \geq 2$$

$$\mathcal{M}_5 = \mathcal{Z}_{21}^{(b_1)} x \mathcal{Z}_{21}^{(b_2)} x \mathcal{Z}_{21}^{(b_3)} x \mathcal{Z}_{21}^{(b_4)} x \mathcal{Z}_{21}^{(b_5)} x \mathcal{D}_{10+|b|} \otimes \mathcal{D}_{10-|b|}; \text{ where } 6 \leq |b| = \sum_{i=1}^5 b_i \leq 10 \text{ and } b_1 \geq 2$$

$$\mathcal{M}_6 = \mathcal{Z}_{21}^{(b_1)} x \mathcal{Z}_{21}^{(b_2)} x \mathcal{Z}_{21}^{(b_3)} x \mathcal{Z}_{21}^{(b_4)} x \mathcal{Z}_{21}^{(b_5)} x \mathcal{Z}_{21}^{(b_6)} x \mathcal{D}_{10+|b|} \otimes \mathcal{D}_{10-|b|}; \text{ where } 7 \leq |b| = \sum_{i=1}^6 b_i \leq 10 \text{ and } b_1 \geq 2$$

$$\mathcal{M}_7 = \mathcal{Z}_{21}^{(b_1)} x \mathcal{Z}_{21}^{(b_2)} x \mathcal{Z}_{21}^{(b_3)} x \mathcal{Z}_{21}^{(b_4)} x \mathcal{Z}_{21}^{(b_5)} x \mathcal{Z}_{21}^{(b_6)} x \mathcal{Z}_{21}^{(b_7)} x \mathcal{D}_{10+|b|} \otimes \mathcal{D}_{10-|b|}; \text{ where } 8 \leq |b| = \sum_{i=1}^7 b_i \leq 10 \text{ and } b_1 \geq 2$$

$$\mathcal{M}_8 = \mathcal{Z}_{21}^{(b_1)} x \mathcal{Z}_{21}^{(b_2)} x \mathcal{Z}_{21}^{(b_3)} x \mathcal{Z}_{21}^{(b_4)} x \mathcal{Z}_{21}^{(b_5)} x \mathcal{Z}_{21}^{(b_6)} x \mathcal{Z}_{21}^{(b_7)} x \mathcal{Z}_{21}^{(b_8)} x \mathcal{D}_{10+|b|} \otimes \mathcal{D}_{10-|b|};$$

$$\text{where } 9 \leq |b| = \sum_{i=1}^8 b_i \leq 10 \text{ and } b_1 \geq 2$$

$$\mathcal{M}_9 = \mathcal{Z}_{21}^{(2)} x \mathcal{Z}_{21}^{(1)} x \mathcal{Z}_{21}^{(1)} x \mathcal{Z}_{21}^{(1)} x \mathcal{Z}_{21}^{(1)} x \mathcal{Z}_{21}^{(1)} x \mathcal{Z}_{21}^{(1)} x \mathcal{Z}_{21}^{(1)} x \mathcal{D}_{20} \otimes \mathcal{D}_0$$

Explore the upshots

The construction and the exactness of the sequence of the contracting homotopies $\{\mathcal{S}_i\}$ where $i = 1, 2, \dots, 8$ discuss in this section as follows

$$\mathcal{S}_0 : \mathcal{M}_0 \rightarrow \mathcal{M}_1, \mathcal{S}_0 \left(\frac{W}{W'} \middle| \frac{1^{(10)} 2^{(\kappa)}}{2^{(10-\kappa)}} \right) = \begin{cases} \mathcal{Z}_{21}^{(\kappa)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(10+\kappa)}}{2^{(10-\kappa)}} \right) & ; \text{ if } \kappa = 2, 3, 4, \dots, 9 \\ 0 & ; \text{ if } \kappa \leq 1 \end{cases},$$

$$\mathcal{S}_1 : \mathcal{M}_1 \rightarrow \mathcal{M}_2, \mathcal{S}_1 \left(\mathcal{Z}_{21}^{(\kappa+1)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+\kappa)} 2^{(\nu)}}{2^{(9-\kappa-\nu)}} \right) \right) = \begin{cases} \mathcal{Z}_{21}^{(\kappa+1)} \chi \mathcal{Z}_{21}^{(\nu)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+\kappa+\nu)}}{2^{(9-\kappa-\nu)}} \right) & ; \text{ if } \nu = 1, 2, \dots, 8 \\ 0 & ; \text{ if } \nu = 0 \end{cases},$$

$$\mathcal{S}_2 : \mathcal{M}_2 \rightarrow \mathcal{M}_3, \mathcal{S}_2 \left(\mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right) \\ = \begin{cases} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\nu)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|+\nu)}}{2^{(9-|\kappa|-\nu)}} \right) & ; \text{ if } \nu = 1, 2, \dots, 7 \\ 0 & ; \text{ if } \nu = 0 \end{cases} \quad ; \text{ where } |\kappa| = \kappa_1 + \kappa_2,$$

$$\mathcal{S}_3 : \mathcal{M}_3 \rightarrow \mathcal{M}_4, \mathcal{S}_3 \left(\mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right) \\ = \begin{cases} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\nu)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|+\nu)}}{2^{(9-|\kappa|-\nu)}} \right) & ; \text{ if } \nu = 1, 2, \dots, 6 \\ 0 & ; \text{ if } \nu = 0 \end{cases} \quad ; \text{ where } |\kappa| = \kappa_1 + \kappa_2 + \kappa_3.$$

$$\mathcal{S}_4 : \mathcal{M}_4 \rightarrow \mathcal{M}_5, \mathcal{S}_4 \left(\mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right) \\ = \begin{cases} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\nu)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|+\nu)}}{2^{(9-|\kappa|-\nu)}} \right) & ; \text{ if } \nu = 1, 2, \dots, 5 \\ 0 & ; \text{ if } \nu = 0 \end{cases} ;$$

where $|\kappa| = \kappa_1 + \kappa_2 + \kappa_3 + \kappa_4$.

$$\mathcal{S}_5 : \mathcal{M}_5 \rightarrow \mathcal{M}_6, \mathcal{S}_5 \left(\mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right) \\ = \begin{cases} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \mathcal{Z}_{21}^{(\nu)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|+\nu)}}{2^{(9-|\kappa|-\nu)}} \right) & ; \text{ if } \nu = 1, 2, \dots, 4 \\ 0 & ; \text{ if } \nu = 0 \end{cases} ;$$

where $|\kappa| = \kappa_1 + \kappa_2 + \kappa_3 + \kappa_4 + \kappa_5$.

$$S_6 : \mathcal{M}_6 \rightarrow \mathcal{M}_7. S_6 \left(Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi Z_{21}^{(K_3)} \chi Z_{21}^{(K_4)} \chi Z_{21}^{(K_5)} \chi Z_{21}^{(K_6)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) \right)$$

$$= \begin{cases} Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi Z_{21}^{(K_3)} \chi Z_{21}^{(K_4)} \chi Z_{21}^{(K_5)} \chi Z_{21}^{(K_6)} \chi Z_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right); & \text{if } v = 1, 2, 3 \\ 0 & ; \text{if } v = 0 \end{cases};$$

where $|K| = K_1 + K_2 + K_3 + K_4 + K_5 + K_6$.

$$S_7 : \mathcal{M}_7 \rightarrow \mathcal{M}_8. S_7 \left(Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi Z_{21}^{(K_3)} \chi Z_{21}^{(K_4)} \chi Z_{21}^{(K_5)} \chi Z_{21}^{(K_6)} \chi Z_{21}^{(K_7)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) \right)$$

$$= \begin{cases} Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi Z_{21}^{(K_3)} \chi Z_{21}^{(K_4)} \chi Z_{21}^{(K_5)} \chi Z_{21}^{(K_6)} \chi Z_{21}^{(K_7)} \chi Z_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right); & \text{if } v = 1, 2 \\ 0 & ; \text{if } v = 0 \end{cases};$$

where $|K| = K_1 + K_2 + K_3 + K_4 + K_5 + K_6 + K_7$.

$$S_8 : \mathcal{M}_8 \rightarrow \mathcal{M}_9. S_8 \left(Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi Z_{21}^{(K_3)} \chi Z_{21}^{(K_4)} \chi Z_{21}^{(K_5)} \chi Z_{21}^{(K_6)} \chi Z_{21}^{(K_7)} \chi Z_{21}^{(K_8)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) \right)$$

$$= \begin{cases} Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi Z_{21}^{(K_3)} \chi Z_{21}^{(K_4)} \chi Z_{21}^{(K_5)} \chi Z_{21}^{(K_6)} \chi Z_{21}^{(K_7)} \chi Z_{21}^{(K_8)} \chi Z_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right); & \text{if } v = 1 \\ 0 & ; \text{if } v = 0 \end{cases};$$

where $|K| = K_1 + K_2 + K_3 + K_4 + K_5 + K_6 + K_7 + K_8$.

Now to prove that $S_i \partial_x + \partial_x S_{i+1} = id_{\mathcal{M}_{i+1}}$, for $i = 0, 1, \dots, 8$.

$$S_0 \partial_x \left(Z_{21}^{(K+1)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+K)} 2^{(V)}}{2^{(9-K)}} \right) \right) = S_0 \partial_{21}^{(K+1)} \left(\frac{W}{W'} \middle| \frac{1^{(11+K)} 2^{(V)}}{2^{(9-K)}} \right) = \binom{K+1+V}{V} Z_{21}^{(K+1+V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+K+V)}}{2^{(9-K-V)}} \right)$$

And

$$\partial_x S_1 \left(Z_{21}^{(K+1)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+K)} 2^{(V)}}{2^{(9-K)}} \right) \right) = \partial_x \left(Z_{21}^{(K+1)} \chi Z_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+K+V)}}{2^{(9-K-V)}} \right) \right)$$

$$= - \binom{K+1+V}{V} Z_{21}^{(K+1+V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+K+V)}}{2^{(9-K-V)}} \right) + Z_{21}^{(K+1)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+K)} 2^{(V)}}{2^{(9-K)}} \right)$$

Clearly $S_0 \partial_x + \partial_x S_1 = id_{\mathcal{M}_1}$.

$$S_1 \partial_x \left(Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) \right)$$

$$= S_1 \left(- \binom{|K|+1}{K_2} Z_{21}^{(|K|+1)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) + Z_{21}^{(K_1+1)} \chi \partial_{21}^{(K_2)} \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) \right)$$

$$= \left(- \binom{|K|+1}{K_2} Z_{21}^{(|K|+1)} \chi Z_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) + \binom{K_2+V}{V} Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2+V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) \right)$$

And

$$\partial_x S_2 \left(Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) \right) = \partial_x \left(Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi Z_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) \right)$$

$$= \binom{|K|+1}{K_2} Z_{21}^{(|K|+1)} \chi Z_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) - \binom{K_2+V}{V} Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2+V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right)$$

$$+ Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right); \text{ where } |K| = K_1 + K_2.$$

Clearly $S_1 \partial_x + \partial_x S_2 = id_{\mathcal{M}_2}$

$$S_2 \partial_x \left(Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2)} \chi Z_{21}^{(K_3)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) \right)$$

$$= S_2 \left[\binom{K_1+1+K_2}{K_2} Z_{21}^{(K_1+1+K_2)} \chi Z_{21}^{(K_3)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) - \binom{K_2+K_3}{K_3} Z_{21}^{(K_1+1)} \chi Z_{21}^{(K_2+K_3)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right) \right]$$

And

Clearly $S_2\partial_x + \partial_x S_3 = id_{\mathcal{M}_3}$.

And

$$\begin{aligned} & \partial_x S_4 \left(Z_{21}^{(\kappa_1+1)} \chi Z_{21}^{(\kappa_2)} \chi Z_{21}^{(\kappa_3)} \chi Z_{21}^{(\kappa_4)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right) \\ &= \partial_x \left(Z_{21}^{(\kappa_1+1)} \chi Z_{21}^{(\kappa_2)} \chi Z_{21}^{(\kappa_3)} \chi Z_{21}^{(\kappa_4)} \chi Z_{21}^{(\nu)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|+\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right) \end{aligned}$$

$$\begin{aligned}
& S_5 \partial_x \left(\mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \mathcal{Z}_{21}^{(\kappa_6)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right) \\
&= S_5 \left[- \binom{(\kappa_1+1+\kappa_2)}{\kappa_2} \mathcal{Z}_{21}^{(\kappa_1+1+\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \mathcal{Z}_{21}^{(\kappa_6)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right. \\
&\quad + \binom{\kappa_2+\kappa_3}{\kappa_3} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2+\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \mathcal{Z}_{21}^{(\kappa_6)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \\
&\quad - \binom{\kappa_3+\kappa_4}{\kappa_4} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3+\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \mathcal{Z}_{21}^{(\kappa_6)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \\
&\quad + \binom{\kappa_4+\kappa_5}{\kappa_5} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4+\kappa_5)} \chi \mathcal{Z}_{21}^{(\kappa_6)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \\
&\quad - \binom{\kappa_5+\kappa_6}{\kappa_6} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5+\kappa_6)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \\
&\quad \left. + \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \partial_{21}^{(\kappa_5)} \mathcal{Z}_{21}^{(\kappa_6)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right] \\
&= - \binom{(\kappa_1+1+\kappa_2)}{\kappa_2} \mathcal{Z}_{21}^{(\kappa_1+1+\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \mathcal{Z}_{21}^{(\kappa_6)} \chi \mathcal{Z}_{21}^{(\nu)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|+\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \\
&\quad + \binom{\kappa_2+\kappa_3}{\kappa_3} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2+\kappa_3)} \chi \mathcal{Z}_{21}^{(\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \mathcal{Z}_{21}^{(\kappa_6)} \chi \mathcal{Z}_{21}^{(\nu)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|+\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \\
&\quad - \binom{\kappa_3+\kappa_4}{\kappa_4} \mathcal{Z}_{21}^{(\kappa_1+1)} \chi \mathcal{Z}_{21}^{(\kappa_2)} \chi \mathcal{Z}_{21}^{(\kappa_3+\kappa_4)} \chi \mathcal{Z}_{21}^{(\kappa_5)} \chi \mathcal{Z}_{21}^{(\kappa_6)} \chi \mathcal{Z}_{21}^{(\nu)} \chi \left(\frac{W'}{W} \middle| \frac{1^{(11+|\kappa|+\nu)}}{2^{(9-|\kappa|-\nu)}} \right)
\end{aligned}$$

And

where $|K| = K_1 + K_2 + K_3 + K_4 + K_5 + K_6$.

$$S_6 \partial_x \left(Z_{21}^{(\kappa_1+1)} \chi Z_{21}^{(\kappa_2)} \chi Z_{21}^{(\kappa_3)} \chi Z_{21}^{(\kappa_4)} \chi Z_{21}^{(\kappa_5)} \chi Z_{21}^{(\kappa_6)} \chi Z_{21}^{(\kappa_7)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right) \\ = S_6 \left[\binom{(\kappa_1 + 1 + \kappa_2)}{\kappa_2} Z_{21}^{(\kappa_1+1+\kappa_2)} \chi Z_{21}^{(\kappa_3)} \chi Z_{21}^{(\kappa_4)} \chi Z_{21}^{(\kappa_5)} \chi Z_{21}^{(\kappa_6)} \chi Z_{21}^{(\kappa_7)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|\kappa|)} 2^{(\nu)}}{2^{(9-|\kappa|-\nu)}} \right) \right]$$

$$\begin{aligned}
&= - \binom{K_1 + 1 + K_2}{K_2} \mathcal{Z}_{21}^{(K_1+1+K_2)} \chi \mathcal{Z}_{21}^{(K_3)} \chi \mathcal{Z}_{21}^{(K_4)} \chi \mathcal{Z}_{21}^{(K_5)} \chi \mathcal{Z}_{21}^{(K_6)} \chi \mathcal{Z}_{21}^{(K_7)} \chi \mathcal{Z}_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1}{2^{(9-|K|-V)}} \right) \\
&+ \binom{K_2 + K_3}{K_3} \mathcal{Z}_{21}^{(K_1+1)} \chi \mathcal{Z}_{21}^{(K_2+K_3)} \chi \mathcal{Z}_{21}^{(K_4)} \chi \mathcal{Z}_{21}^{(K_5)} \chi \mathcal{Z}_{21}^{(K_6)} \chi \mathcal{Z}_{21}^{(K_7)} \chi \mathcal{Z}_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) \\
&- \binom{K_3 + K_4}{K_4} \mathcal{Z}_{21}^{(K_1+1)} \chi \mathcal{Z}_{21}^{(K_2)} \chi \mathcal{Z}_{21}^{(K_3+K_4)} \chi \mathcal{Z}_{21}^{(K_5)} \chi \mathcal{Z}_{21}^{(K_6)} \chi \mathcal{Z}_{21}^{(K_7)} \chi \mathcal{Z}_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) \\
&+ \binom{K_4 + K_5}{K_5} \mathcal{Z}_{21}^{(K_1+1)} \chi \mathcal{Z}_{21}^{(K_2)} \chi \mathcal{Z}_{21}^{(K_3)} \chi \mathcal{Z}_{21}^{(K_4+K_5)} \chi \mathcal{Z}_{21}^{(K_6)} \chi \mathcal{Z}_{21}^{(K_7)} \chi \mathcal{Z}_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) \\
&- \binom{K_5 + K_6}{K_6} \mathcal{Z}_{21}^{(K_1+1)} \chi \mathcal{Z}_{21}^{(K_2)} \chi \mathcal{Z}_{21}^{(K_3)} \chi \mathcal{Z}_{21}^{(K_4)} \chi \mathcal{Z}_{21}^{(K_5+K_6)} \chi \mathcal{Z}_{21}^{(K_7)} \chi \mathcal{Z}_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) \\
&- \binom{K_6 + K_7}{K_7} \mathcal{Z}_{21}^{(K_1+1)} \chi \mathcal{Z}_{21}^{(K_2)} \chi \mathcal{Z}_{21}^{(K_3)} \chi \mathcal{Z}_{21}^{(K_4)} \chi \mathcal{Z}_{21}^{(K_5)} \chi \mathcal{Z}_{21}^{(K_6+K_7)} \chi \mathcal{Z}_{21}^{(V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) \\
&- \binom{K_7 + V}{V} \mathcal{Z}_{21}^{(K_1+1)} \chi \mathcal{Z}_{21}^{(K_2)} \chi \mathcal{Z}_{21}^{(K_3)} \chi \mathcal{Z}_{21}^{(K_4)} \chi \mathcal{Z}_{21}^{(K_5)} \chi \mathcal{Z}_{21}^{(K_6)} \chi \mathcal{Z}_{21}^{(K_7+V)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|+V)}}{2^{(9-|K|-V)}} \right) \\
&+ \mathcal{Z}_{21}^{(K_1+1)} \chi \mathcal{Z}_{21}^{(K_2)} \chi \mathcal{Z}_{21}^{(K_3)} \chi \mathcal{Z}_{21}^{(K_4)} \chi \mathcal{Z}_{21}^{(K_5)} \chi \mathcal{Z}_{21}^{(K_6)} \chi \mathcal{Z}_{21}^{(K_7)} \chi \left(\frac{W}{W'} \middle| \frac{1^{(11+|K|)} 2^{(V)}}{2^{(9-|K|-V)}} \right);
\end{aligned}$$

where $|K| = K_1 + K_2 + K_3 + K_4 + K_5 + K_6 + K_7$.

Clearly $S_6 \partial_x + \partial_x S_7 = \text{id}_{\mathcal{M}_7}$.

[illegible]

And

where $|K| = K_1 + K_2 + K_3 + K_4 + K_5 + K_6 + K_7 + K_8$.

Clearly $S_7\partial_x + \partial_x S_8 = id_{\mathcal{M}_8}$

Thus $\{S_i\}$; where $i = 0, 1, 2, \dots, 8$ is a contracting homotopy¹⁰ which implies the complex

$$0 \rightarrow \mathcal{M}_9 \rightarrow \mathcal{M}_8 \rightarrow \mathcal{M}_7 \rightarrow \mathcal{M}_6 \rightarrow \mathcal{M}_5 \rightarrow \mathcal{M}_4 \rightarrow \mathcal{M}_3 \rightarrow \mathcal{M}_2 \rightarrow \mathcal{M}_1 \rightarrow \mathcal{M}_0$$

is exact.

Results and discussion

In this article, the terms of the characteristic-free resolution for the situation (11,10; 1) and the exactness of the Weyl resolution of the sequence are surveyed after obtaining the contracting homotopy of $\{S_i\}$; where $i = 0, 1, \dots, 8$.

To get the results of this work the important computations which are $S_{i-1}\partial_x + \partial_x S_i = id_{\mathcal{M}_i}$; where $i = 0, 1, 2, \dots, 8$ proved.

Conclusion

The exact complex sequence of the terms of the characteristic-free resolution got in this work from the contracting homotopies of $\{S_i\}$ where $i = 0, 1, 2, \dots, 8$ for the case (11,10;1) after finding all the terms of the Weyl module and the Weyl module resolution, also by using the form of the letter place the maps of $\{S_i\}$ where $i = 0, 1, 2, \dots, 8$ are described and $\partial_{21}^{(e)}$ the composition of place polarizations from places 1 and 2 of degree e to the negative place $\{1', 2', \dots, 12'\}$ act on. $\mathcal{D}_{10+e} \oplus \mathcal{D}_{10-e}$.

Acknowledgment

Our thanks and gratitude extend to every member in Baghdad Science journal, College of Education for Pure Science Ibn-Al-Haitham, University of Baghdad and College of Computer Science and Mathematics, University of Thi-Qar for their support, encouragement and cooperation they provided to accomplish this research.

Authors' declaration

- Conflicts of Interest: None
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been included with the necessary permission for re-publication, which is attached to the manuscript.
- No animal studies are present in the manuscript.
- No human studies are present in the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee at University of Baghdad.

Authors' contribution statement

N. S. J. presented the idea and discussed it theoretically and checked the computations. W. K. J. made the computations for this work, N. S.J. and W. K. J. discussed and validates the results obtained by manual calculations.

References

1. Haytham RH, Alaa OA. Complex of lascaroux in partition (3,3,2). Ibn al-Haitham J Pure Appl. Sci. 2015;28(1):171–178.
2. Khalaf SA, Jasim NS. Calculation for the groups $SL(2, \mathfrak{A})$, $\mathfrak{A} = 31$ and 37. J Discrete Math Sci Cryptogr. 2022;25(2):609–613. <https://doi.org/10.1080/09720529.2021.1972614>.
3. Hassan HR. The reduction of resolution of weyl module from characteristic-free resolution in case (4,4,3). Ibn al-Haitham J Pure Appl Sci. 2012;25(3):341–355.

4. Hameed MI, Shihab BN. On differential subordination and superordination for univalent function involving new operator. I.J.C.S.M. 2022;3(1):22–31. <https://doi.org/10.52866/ijcsm.2022.01.01.003>.
5. Hamel MA, Khalaf HY. Fuzzy maximal sub-modules. Iraqi J Sci. 2020;61(5):164–1172. <https://doi.org/10.24996/ijs.2020.61.5.24>.
6. Aljebory KS, Alhossaini AMA. Almost projective semimodules. Baghdad Sci J. 2023;20(6):2271–2277. <https://doi.org/10.21123/bsj.2023.7716>.
7. Farhan E. On semigroup ideals and right n-derivation in 3-prime near-rings. Baghdad Sci J. 2024;21(1):196–203. <https://doi.org/10.21123/bsj.2023.8086>.
8. Ali SH, Kareem FF, Hameed AT. Hyper Fuzzy AT-ideals of AT-algebra. International Conference of Modern Applications on Information and Communication Technology (ICMAICT) 22–23 October 2020, University of Babylon, Babylon-Hilla City, Iraq. J Phys.: Conf Ser. 2021;1804(1):1–15.
9. Ahmed G, Gedaan S, Ibrahim R. Spectrum of Secondary Submodules. Ibn Al-Haitham International Conference for Pure and Applied Sciences (IHICPS) 9–10 December 2020, Baghdad, Iraq. J Phys.: Conf Ser. 2021;1879(3):1–20. <https://doi.org/10.1088/1742-6596/1879/3/032007>.
10. Buchsbaum DA. A characteristic-free example of Lascoux resolution and letter place methods for intertwining numbers. Eur J Comb. 2004;25:1169–1179. <https://doi.org/10.1016/j.ejc.2003.09.017>.

نتيجة للحالة (11,10; 1)

وليد خالد جابر^{1,2*}، نيران صباح جاسم¹

¹قسم الرياضيات، كلية التربية للعلوم الصرفة (ابن الهيثم)، جامعة بغداد، بغداد، العراق.

²قسم الرياضيات، كلية علوم الحاسوب والرياضيات، جامعة ذي قار، ناصرية، العراق.

الخلاصة

خلال هذا العمل R هي حلقة ابدالية ذات 1، $\mathcal{F}$ هو مقياس R الحر و $\mathcal{D}_i$ هو جبر قوى القسمة من الدرجة i . التجزئة ذات الطول $\ell(\lambda)$ $= n$ هي السلسلة $(\lambda_1, \lambda_2, \dots, \lambda_n)$ من الصحيحات غير السالبة وبترتيب غير متزايد $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > 0$. وزن التجزئة λ هو $|\lambda| = \lambda_1 + \lambda_2 + \dots + \lambda_n$. المتسلسلة النسبية هي الزوج (λ, μ) من المتسلسلات بحيث ان $\mu \leq \lambda$ ويرمز لها بالرمز λ/μ . اذا كان كلا من λ و μ تجزئة فان السلسلة النسبية λ/μ تسمى تجزئة منحرفة. ليكن المولد الحر لجبر قوى القسمة $\mathcal{Z}_{21}$ بمولد واحد. عنصر قوى القسمة $\mathcal{Z}_{21}^{(e)}$ من الدرجة e للمولد الحر $\mathcal{Z}_{21}$ يمثل في $\mathcal{D}_{(p+e)} + \mathcal{D}_{(q-e)}$ بالاستقطاب المكاني من الدرجة e من المكان 1 الى المكان 2. التدرج الجبري $\mathcal{A} = \mathcal{D}(\mathcal{Z}_{21})$ يمثل بتدرج المقياس $\mathcal{M}_{(q-e)}$ ، $\mathcal{M} = \mathcal{D}_{(p+e)} + \mathcal{D}_{(q-e)} = \Sigma \mathcal{M}_{(q-e)}$ ، درجة المقسوم الثاني تعتبر التدرج. M هو مقياس $A =$ الايسر المتدرج، حيث لكل $w = \mathcal{Z}_{21}^{(e)} \in \mathcal{A}$ و $v \in \mathcal{D}_{(p+e)} + \mathcal{D}_{(q-e)}$ فانه لدينا $w(v) = \mathcal{Z}_{21}^{(e)}(v) = \partial_{21}^{(e)}(v)$. البحث يدرس تمام تحليل وايل للمتسلسلة للحالة (11,10; 1) بعد الحصول على التعاقد الهوموتوبي لـ $\{S_i\}$ ؛ حيث $i = 0, 1, 2, \dots, 8$ ، و عناصر تحليل المميز - الحر لها.

الكلمات المفتاحية: جبر قوى القسمة، موقع الحرف، تداخل، موقع الاستقطابات، تحليل، مقياس وايل.